



# Aminogro<sup>®</sup>

## 8% Boron

### MAXi

Organic Carbon: 14% w/w Amine Nitrogen: 6.4% w/w

**OrganoPLEX<sup>®</sup>** L-form Amino Acid  
Chelate Carboxylate Mannitol Complex

#### PLANT FUNCTION

- Cell wall formation & elasticity • Sugar transportation • Helps calcium translocation
- Flower retention, pollen formation & germination • Assists with hormone production

#### DEFICIENCY SYMPTOMS

- Appear at growing points • Stunted plants • Dead or misshapen young leaves
- Red pigmentation in leaves • Hollow stems & fruit • Low seed production
- Flower, bud or fruit drop



**ORGANIC CROP PROTECTANTS**  
1956 Dandenong Rd, Clayton VIC 3168  
Telephone: 1800 634 204 (Free Call)  
ocp.com.au



**CONTENTS**  
**25 kg**

Y20-3312

#### GENERAL INFORMATION

Aminogro<sup>®</sup> MAXi Boron provides plants with a boron chelate using L-form free amino acids and other simple organic sugars and acids. Nutrients are absorbed and shuttled around inside plants in the form of amino acids, simple sugars and acids so OrganoPLEX<sup>™</sup> is an efficient way of delivering and mobilizing boron in plants. The organic carbon in Aminogro<sup>®</sup> MAXi Boron also enhances microbial activity in soil which assists the plant accessing boron and other nutrients when applied to the soil.

#### DIRECTIONS FOR USE:

The rates below are a guide and dependent upon soil and sap tests throughout the growing season. Foliar applications of boron are more effective than soil applications. Consult your OCP Agronomist for further advice.

Tree Crop	Rate	Application Timing
Pome Fruit	1-2 kg/ha	1. 1-2 weeks before flowering (if better fruit set is desired) 2. 1 month after fruit set 3. 2 months before harvest 4. 2 weeks before harvest
Stone Fruit	1-2 kg/ha	1. 1 week before flowering (if better fruit set is desired) 2. 2-3 weeks later 3. 2-3 weeks later 4. 2 weeks before harvest
Nuts	1.5-2.5 kg/ha	1. 2-3 weeks after nut set 2. After picking to extend leaf function & sugar accumulation
Bananas	1-2 kg/ha	With seasonal applications of potassium & nitrogen
Vine Crop	Rate	Application Timing
Table Grapes	1-2 kg/ha	1. 1 week before flowering 2. 4 weeks later 3. 2 weeks before harvest 4. Post harvest
Wine Grapes	1-2 kg/ha	1. 1-2 weeks before flowering 2. Veraison 3. Post harvest
Strawberries/Blueberries	1-2 kg/ha	Begin just before flowering & repeat as sap results indicate
Protected Crops	Rate	Application Timing
Hydroponic run to waste & NFT crops	200 g/10,000 L	Consult OCP agronomist for crop specific requirements
Nursery Stock	Foliar	If deficiency is detected consult OCP agronomist

Field Crops	Rate	Application Timing
Cereal, Soybeans	2-3 kg/ha	As required by soil test results
Leafy Veggies Lettuce	1-2 kg/ha	1. 4 leaf stage 2. Every 14 days until harvest
Carrots, Onions, Leeks, Turnips	2 kg/ha	1. 2 weeks after emergence 2. At root enlargement 3. 6 weeks before harvest
Brassica, Crucifer	2 kg/ha	1. 6-8 true leaf stage 2. 10-14 days later 3. At head initiation
Bean & Peas	2 kg/ha	1. 4-6 true leaf stage 2. At first bloom 3. At first pod
Potatoes, Yam, Taro	2 kg/ha	1. (Seed treatment optional) 2. At tuber set 3. Early bloom
Sweet Corn	2 kg/ha	1. 2-6 true leaf stage 2. 50-75cm growth 3. Just prior to tasselling
Tomato, Capsicum	2 kg/ha	1. 15-20 cm 2. First pre-bloom 3. 7 days later
Cucurbits	2 kg/ha	1. 4 true leaf stage 2. Pre-bloom 3. First fruit set
Professional Turf	Rate	Application Timing
Greens & Tees	200 g/1000 m <sup>2</sup>	As required by soil test
Fairways	1 kg/ha	As required by soil test

#### COMPATIBILITY

Aminogro<sup>®</sup> MAXi Boron can be mixed with most horticultural sprays. However, the performance of some sprays may be affected by high cation levels like calcium, potassium and phosphorus. It is best to apply these sprays separately or conduct a jar test to determine compatibility in the ratios they are to be used. Always check pesticide labels for compatibility with foliar fertilisers, or test the mixture on a small area before widespread application.

For further information contact Organic Crop Protectants  
Free Call: 1800 634 204 Email: info@ocp.com.au  
**FOR EMERGENCIES ONLY, CALL AUSTRALIA 1800 033 111 (ALL HOURS)**

Aminogro and the OrganoPLEX device are registered trade marks and the OCP drop device is a trade mark of Organic Crop Protectants



Y20-3320



Job OCP572 AminogroMaxi BORON 25kg

Version

Size 208mm x 190mm  
Colours



Barcode % 100%  
Spot



Process  
Flexo  
Lytho  
Gravure  
Offset

**FA**

05/05/20