



Aminogro[®] MAXi

N13

Organic Nitrogen 13% w/w

Aminogro[®] MAXi N13 provides plants with an array of amino acids in the form of L-form free amino acids, peptides and polypeptides. Nitrogen is carried around inside plants in the form of amino acids so fertilising with amino acids is an efficient way of delivering nitrogen to the plant. The organic carbon in Aminogro[®] MAXi N13 also enhances microbial activity which assists the plant to access other nutrients. Aminogro[®] MAXi N13 also acts as a natural chelator of other nutrients in tank mixes.



ORGANIC CROP PROTECTANTS
1956 Dandenong Rd, Clayton VIC 3168
Telephone: 1800 634 204 (Free Call)
ocp.com.au



CONTENTS
25 kg

Y20-3100

GENERAL INFORMATION

The rates below are a guide and dependent upon the amount of nitrogen applied as a basal fertiliser pre-season and crop yield targets. Adjustments should be made to nitrogen requirements throughout the growing season based on sap tests. Consult your OCP Agronomist for further advice.

Tree Crop	Rate	Application Timing
Pome Fruit	500 g/100 L or max 5 kg/ha	1. Green tip to tight cluster 2. Open cluster to pink 3. 1st cover to 4th cover 4. Every 2-3 wks after 1st cover 5. 2-3 wks before harvest
Stone Fruit	500 g/100 L or max 5 kg/ha	1. At petal fall 2. 10-14 days later 3. Post harvest- drip x 2 apps 10kg/ha
Nuts	500 g/100 L or max 5 kg/ha	1. At petal fall 2. 10-14 days later 3. Post harvest- drip x 2 apps 10kg/ha
Bananas	1 kg/100 L or max 50 kg/ha	With every spray or as required through fertigation
Vine Crop	Rate	Application Timing
Table Grapes	500 g/100 L or max 5 kg/ha	1. 10-15 cm canes 2. 25-35 cm canes 3. Berry set 4. 7 days later 5. 7 days later Post harvest- drip x 2 apps
Wine Grapes	300 g/100 L or max 3 kg/ha	1. Berry set 2. Veraison* 3. Pre-bloom* 4. Berry set* 5. Veraison* Post harvest- drip x 2 apps 10kg/ha * For growers who want bigger berries and better set
Strawberries/ Blueberries	5 kg/ha	Begin at fruit set and repeat as sap results indicate

Field Crops	Rate	Application Timing
Cereal, Soybeans	2-3 kg/ha	As required by soil or sap test results.
Lettuce	500 g/100 L or max 10 kg/ha	1. Seed treatment 2. 4 leaf stage 3. Every 14 days until harvest
Carrots, Onions, Leeks, Turnips	500 g/100 L or max 10 kg/ha	1. 2 weeks after emergence 2. At root enlargement 3. 10-14 days until harvest
Brassica, Crucifer	500 g/100 L or max 10 kg/ha	1. 4-6 true leaf stage 2. 10-14 days later 3. At head initiation
Bean & Peas	500 g/100 L or max 10 kg/ha	1. 4-6 true leaf stage 2. At first bloom 3. At first pod
Potatoes, Yam, Taro	500 g/100 L or max 10 kg/ha	1. (Seed treatment optional) 2. At tuber set 3. 10-14 days later 4. Early bloom
Sweet Corn	500 g/100 L or max 10 kg/ha	1. 2-6 true leaf stage 2. 50-75cm growth 3. Just prior to tasselling
Protected Crops	Rate	Application Timing
Hydroponic NFT All Crops	5 kg/10,000 L	Every two weeks. Consult OCP agronomist if using the product for entire nitrogen source
Cutflowers	Drip & Foliar	Every 2nd wk @ 2.5kg / 2000 plants as a drip and 250 g / 100L as foliar every wk
Nursery Stock	Foliar	Every week @ 500g / 100L

Professional turf	Rate	Application timing	General Information
Greens, Tee & Intensively Managed Sports Turf	5 kg/ha	Monthly applications or as required for aesthetic appeal	Aminogro [®] MAXi Iron is compatible with PGR's. If mixing in alkaline water, adjust to pH 6-6.5 using a suitable buffer
Fairways & Sports Fields	2 kg/ha	Monthly application for optimum greening	

COMPATIBILITY

Aminogro[®] MAXi N13 can be mixed with most horticultural sprays. Always check pesticide labels for compatibility with foliar fertilisers, or test the mixture on a small area before widespread application.

For further information contact Organic Crop Protectants
Free Call: 1800 634 204 Email: info@ocp.com.au
FOR EMERGENCIES ONLY, CALL AUSTRALIA 1800 033 111 (ALL HOURS)

Aminogro is a registered trade mark and the OCP drop device is a trade mark of Organic Crop Protectants.



Y20-3101



Job OCP572 AminogroMaxi N13 25kg	Version		
Size 208mm x 190mm	Barcode % 100%	v1	
Colours	Spot	24/06/20	
C M Y K	○ ○ ○ ○		
		Process Flexo Lytho Gravure Offset	